

## Module 3.7

### Numerical Integration and the Lognormal Distribution

1

## Overview

- Explore numerical integration
  - Lognormal distribution
  - Normal distribution
  - Deep dive into properties as applied in finance
- Numerous comparisons and contrasts
- Bivariate distributions



© Financial Risk Management, LLC

2

## Univariate Distributions

- Cumulative distribution function (CDF)

$$F_X(x) \equiv \Pr(X \leq x) \text{ (generic)}$$

$$F_X(x) = \int_{-\infty}^x f_X(j) dj \text{ (continuous)}$$

- Probability density function (continuous)

$$f_X(j)$$



© Financial Risk Management, LLC

3

## Lognormal Distribution

- Parameters:  $\mu$  (Mean),  $\sigma$  (Std. Dev.)

- Limits

- Mean:  $-\infty < \mu < +\infty$

- Standard deviation:  $\sigma > 0$

- Range:  $0 < x < +\infty$

- Note:  $x$  cannot equal zero

- If  $y = \ln(x)$  is normally distributed (ND), then  $x$  is said to have a lognormal distribution (LD)



© Financial Risk Management, LLC

4

## ND and LD CDF and PDF

- CDF:  $\Lambda(\mu, \sigma) = \Lambda(d) = \int_{-\infty}^d \frac{e^{-\frac{[\ln(x)-\mu]^2}{2\sigma^2}}}{x\sigma\sqrt{2\pi}} dx$  (Lognormal CDF) and

$$N(\mu, \sigma) = N(d) = \int_{-\infty}^d \frac{e^{-\frac{(x-\mu)^2}{2\sigma^2}}}{\sigma\sqrt{2\pi}} dx \text{ (Normal CDF).}$$

- PDF:

$$\lambda(\mu, \sigma) = \lambda(x) = \frac{e^{-\frac{[\ln(x)-\mu]^2}{2\sigma^2}}}{x\sigma\sqrt{2\pi}} \text{ (Lognormal PDF) and}$$

$$n(\mu, \sigma) = n(x) = \frac{e^{-\frac{(x-\mu)^2}{2\sigma^2}}}{\sigma\sqrt{2\pi}} \text{ (Normal PDF).}$$



© Financial Risk Management, LLC

5

## Median, Mode, Mean

- Median:  $\hat{x}$  such that  $\int_0^{\hat{x}} f(x) dx = \int_{\hat{x}}^{\infty} f(x) dx = \frac{1}{2}$ .

$$\text{Median}_\lambda = e^\mu \text{ and}$$

$$\text{Median}_n = \mu.$$

- Mode:  $f'(x) = 0$  and  $f''(x) < 0$   $\text{Mode}_\lambda = e^{\mu - \sigma^2}$  and  $\text{Mode}_n = \mu$ .

- Mean:  $\text{Mean}_\lambda = e^{\mu + \frac{\sigma^2}{2}}$  and  $\text{Mean}_n = \mu$ .



© Financial Risk Management, LLC

6

## Variance and Skewness

- Variance:  $\mu_{2\lambda} = \text{Variance}_\lambda = e^{2\mu + \sigma^2} (e^{\sigma^2} - 1)$  and  $\mu_{2n} = \text{Variance}_n = \sigma^2$ .
- Skewness:  $\mu_{3\lambda} = e^{3\mu + \frac{3\sigma^2}{2}} (e^{\sigma^2} - 1)^2 (e^{\sigma^2} + 2)$  and  $\mu_{3n} = 0$ .
- Normalized Skewness:  $\gamma_{S\lambda} = \frac{\mu_{3\lambda}}{\mu_{2\lambda}^{3/2}} = \sqrt{e^{\sigma^2} - 1} (e^{\sigma^2} + 2)$  and  $\gamma_{Sn} = \frac{\mu_{3n}}{\mu_{2n}^{3/2}} = 0$ .



© Financial Risk Management, LLC

7

7

## Kurtosis

- Kurtosis:  $\mu_{4\lambda} = e^{4\mu + 2\sigma^2} (e^{\sigma^2} - 1)^2 (e^{4\sigma^2} + 2e^{3\sigma^2} + 3e^{2\sigma^2} - 3)$  and  $\mu_{4n} = 3\sigma^4$ .
- Excess kurtosis:  $\gamma_{K\lambda} = \frac{\mu_{4\lambda}}{\mu_{2\lambda}^2} - 3 = e^{4\sigma^2} + 2e^{3\sigma^2} + 3e^{2\sigma^2} - 6$  and  $\gamma_{Kn} = \frac{\mu_{4n}}{\mu_{2n}^2} - 3 = 0$ .



© Financial Risk Management, LLC

8

8

## Asset Price Distributions

- If  $x \sim N(\mu_g, \sigma_g)$  (g denotes generic) and  $y = e^x$ , then  $y \sim \Lambda(\mu_g, \sigma_g)$
- Rates of return:  $S_t = S_0 e^{R(T-t)}$
- Distribution of returns:  $R \sim N[\mu(T-t), \sigma\sqrt{T-t}]$
- Distribution of prices:  $S_T \sim \Lambda[\ln(S_0) + \mu(T-t), \sigma\sqrt{T-t}]$



© Financial Risk Management, LLC

9

9

## Asset Price Distribution

- Terminal stock prices:  $E(S_T) = S_0 e^{\left(\frac{\mu + \sigma^2}{2}\right)(T-t)}$   $Var(S_T) = S_0^2 \left( e^{2(\mu + \sigma^2)(T-t)} - e^{2\mu + \sigma^2(T-t)} \right)$
- Returns:  $\mu = \ln\left[\frac{E(S_T)}{S_0}\right] / (T-t)$   $\sigma^2 = \ln\left[\frac{Var(S_T)}{E(S_T)^2} + 1\right] / (T-t)$



© Financial Risk Management, LLC

10

10

## Asset Price Distributions

- Suppose  $S = 100$ ,  $T = 1$ ,  $\hat{\mu} = 12$ ,  $\sigma = 30\%$
- Intend:  $E(S_T) = S_0 e^{\hat{\mu}(T-t)} = 100e^{0.12(1)} = 112.749685$
- Two versions:  $E(S_T) = S_0 e^{\left(\frac{\mu + \sigma^2}{2}\right)(T-t)}$  and  $E(S_T) = S_0 e^{\left(\frac{\mu + \sigma^2}{2}\right)(T-t)}$

$$E(S_T) = S_0 e^{\left(\frac{\mu + \sigma^2}{2}\right)(T-t)} = S_0 e^{\hat{\mu}(T-t)} = 100e^{0.12(1)} = 112.749685$$

$$\mu = \hat{\mu} - \frac{\sigma^2}{2} = 0.12 - \frac{0.3^2}{2} = 0.075$$



© Financial Risk Management, LLC

11

11

## Asset Price Distributions

$$\begin{aligned} S_T &\sim \Lambda\left[\ln(S_0) + \left(\hat{\mu} - \frac{\sigma^2}{2}\right)(T-t), \sigma\sqrt{T-t}\right] \\ &= \Lambda\left[\ln(S_0) + \mu(T-t), \sigma\sqrt{T-t}\right] \\ &= \Lambda\left[\ln(100) + 0.075(1), 0.30\sqrt{1}\right] = \Lambda(4.680170, 0.30) \\ \text{Median} &= e^{\ln(S_0) + \left(\hat{\mu} - \frac{\sigma^2}{2}\right)(T-t)} = e^{4.680170} = 107.788415, \\ \text{Mode} &= e^{\ln(S_0) + \left(\hat{\mu} - \frac{\sigma^2}{2}\right)(T-t) - \sigma^2(T-t)} = e^{4.680170 - 0.09(1)}, \text{ and} \\ &= e^{4.680170 - 0.09(1)} = 98.511194 \\ \text{Variance} &= e^{2\left[\ln(S_0) + \left(\hat{\mu} - \frac{\sigma^2}{2}\right)(T-t)\right] + \sigma^2(T-t)} - e^{2\left[\ln(S_0) + \left(\hat{\mu} - \frac{\sigma^2}{2}\right)(T-t)\right]} \\ &= e^{2\left[4.680170 + 0.09(1)\right] + 0.09(1)} - e^{2\left[4.680170 + 0.09(1)\right]} = 1,197.189781. \end{aligned}$$



standard deviation is 34.600430.

© Financial Risk Management, LLC

12

12

## Role of Standard Deviation

- Increasing return standard deviation (volatility) implies terminal price has
  - Increasing mean
  - Constant median
  - Decreasing mode
  - Increasing skewness
  - Increasing likelihood of very low values



© Financial Risk Management, LLC

13

13

Figure 3.7.1A. Stock Price = \$100, Horizon = 1 year Expected Return = 12%, Standard Deviation = 30%




14

Figure 3.7.1B. Stock Price = \$100, Horizon = 1 year Expected Return = 12%, Standard Deviation = 30%




15

Figure 3.7.2A. Stock Price = \$100, Horizon = 1 year Expected Return = 12%, Standard Deviation = 80%




16

Figure 3.7.2B. Stock Price = \$100, Horizon = 1 year Expected Return = 12%, Standard Deviation = 80%




17


Figure 3.7.3A. Stock Price = \$100, Horizon = 1 year Expected Return = 12%, Standard Deviation = 130%




18



19



20



21

### Probability Call In-The-Money

- Terminal stock price:  $\tilde{S}_T = S_0 e^{\left(\mu - \frac{\sigma^2}{2}\right)T + \sigma\sqrt{T}\tilde{\varepsilon}}$
- Probability:  $\Pr(S_T > X) = \Pr\left[S_0 e^{\left(\mu - \frac{\sigma^2}{2}\right)T + \sigma\sqrt{T}\tilde{\varepsilon}} > X\right]$

$$\Pr(S_T > X) = \Pr\left[\ln\left(\frac{S_0}{X}\right) + \left(\mu - \frac{\sigma^2}{2}\right)T + \sigma\sqrt{T}\tilde{\varepsilon} > 0\right]$$

$$= \Pr\left[\tilde{\varepsilon} > \frac{-\ln\left(\frac{S_0}{X}\right) - \left(\mu - \frac{\sigma^2}{2}\right)T}{\sigma\sqrt{T}}\right] = 1 - \Pr\left[\tilde{\varepsilon} < -\frac{\ln\left(\frac{S_0}{X}\right) + \left(\mu - \frac{\sigma^2}{2}\right)T}{\sigma\sqrt{T}}\right]$$

$$= 1 - N(-d_1) = N(d_1)$$

© Financial Risk Management, LLC 22

22

### Multivariate Distributions

- Finance suffers from “curse of dimensionality”
- S&P 500 index
  - Technically a 500 dimensional problem (each stock)
  - Each stock has multiple factors or dimensions
  - Quants aggressively seek to reduce dimensionality
  - In practice, one rarely gets the dimensions down to only one. Hence, we need the capacity to model multiple dimensional problems.

© Financial Risk Management, LLC 23

23

### Multivariate Normal

- PDF:  $n_N(\mathbf{x}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{e^{-\frac{(\mathbf{x}-\boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}{2}}}{\sqrt{(2\pi)^N |\boldsymbol{\Sigma}|}}$
- If  $\mathbf{x} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  and  $\mathbf{y} = e^{\mathbf{x}}$ , then
  - $E(y_i) = e^{\mu_i + \frac{\sigma_i^2}{2}}$  and
  - $\text{cov}(y_i, y_j) = e^{\mu_i + \mu_j + \frac{1}{2}(\sigma_i^2 + \sigma_j^2)}(e^{\Sigma_{ij}} - 1)$ .

© Financial Risk Management, LLC 24

24

## Bivariate PDFs and CDFs

### Normal:

$$n_2(x_1, x_2; \rho) \equiv \frac{e^{-\frac{\left(\frac{x_1 - \mu_1}{\sigma_1}\right)^2 - 2\rho\left(\frac{x_1 - \mu_1}{\sigma_1}\right)\left(\frac{x_2 - \mu_2}{\sigma_2}\right) + \left(\frac{x_2 - \mu_2}{\sigma_2}\right)^2}{2(1 - \rho^2)}}{2\pi\sigma_1\sigma_2\sqrt{1 - \rho^2}}$$

$$N_2(a, b; \rho) \equiv \int_{-\infty}^a \int_{-\infty}^b n_2(x_1, x_2; \rho) dx_1 dx_2$$

### Lognormal:

$$\lambda_2(x_1, x_2; \rho) \equiv \frac{e^{-\frac{\left[\frac{\ln(x_1) - \mu_1}{\sigma_1}\right]^2 - 2\rho\left[\frac{\ln(x_1) - \mu_1}{\sigma_1}\right]\left[\frac{\ln(x_2) - \mu_2}{\sigma_2}\right] + \left[\frac{\ln(x_2) - \mu_2}{\sigma_2}\right]^2}{2(1 - \rho^2)}}}{2\pi x_1 x_2 \sigma_1 \sigma_2 \sqrt{1 - \rho^2}}$$


$$\Lambda_2(a, b; \rho) \equiv \int_{-\infty}^a \int_{-\infty}^b \lambda_2(x_1, x_2; \rho) dx_1 dx_2$$



25

## Bivariate Lognormal Simulation

Figure 3.7.5 Bivariate Lognormal Simulation



26

## Conditional Normal Distribution

### Note

$$n(x_2 | x_1) = \frac{e^{-\frac{\left[x_2 - \left[\mu_2 + \rho\frac{\sigma_2}{\sigma_1}(x_1 - \mu_1)\right]\right]^2}{2\sigma_2^2(1 - \rho^2)}}}{\sigma_2\sqrt{2\pi(1 - \rho^2)}}$$

### Thus, $n(x_2 | x_1) \sim N(\mu_{2|1}, \sigma_{2|1}^2)$

$$\mu_{2|1} = \mu_2 + \rho\frac{\sigma_2}{\sigma_1}(x_1 - \mu_1) \text{ and}$$

$$\sigma_{2|1}^2 = \sigma_2^2(1 - \rho^2)$$



27

## Contrasting ND and LD


### Next several graphs generated in R illustrate role of volatility on

- PDFs
- CDFs
- Mean, Median, and Mode
- Simulations




28

Figure 3.7.6 Univariate Normal PDFs (30%, 80%, 130%, 180%)




29


Figure 3.7.7 Univariate Lognormal PDFs (\$34, \$106.75, \$237, \$558)




30




31



32



33



34

**Summary**

- Explored numerical integration
  - Lognormal distribution
  - Normal distribution
  - Deep dive into properties as applied in finance
- Numerous comparisons and contrasts
- Bivariate distributions

© Financial Risk Management, LLC

35